



## **Sensitivity of a satellite-derived drought index under soil moisture-limited vs. energy-limited evapotranspiration.**

**Garcia, Monica; Fernandez, Nestor; Villagarcia, Luis; Domingo, Francisco; Puigdefabregas, Juan; Sandholt, Inge**

*Publication date:*  
2014

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Garcia, M., Fernandez, N., Villagarcia, L., Domingo, F., Puigdefabregas, J., & Sandholt, I. (2014). *Sensitivity of a satellite-derived drought index under soil moisture-limited vs. energy-limited evapotranspiration..* Poster session presented at 7th International Scientific Conference on the Global Water and Energy Cycle, Hague, Netherlands. [http://gewex.org/2014conf/pdfs/Garcia\\_7-6.pdf](http://gewex.org/2014conf/pdfs/Garcia_7-6.pdf)

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.





# Sensitivity of a satellite-derived drought index under soil moisture-limited vs. energy-limited evapotranspiration

Mónica García <sup>(1)</sup>, Nestor Fernández<sup>(2)</sup>, Luis Villagarcía<sup>(3)</sup>, Francisco Domingo<sup>(4)</sup>, Juan Puigdefábregas<sup>(4)</sup>, Inge Sandholt<sup>(1)</sup>

<sup>(1)</sup> Denmark Technical University (DTU), 2800 Kgs. Lyngby, Denmark. E-mail: [mgarc@env.dtu.dk](mailto:mgarc@env.dtu.dk),

<sup>(2)</sup> Department of Conservation Biology. Estación Biológica de Doñana, CSIC, 41092 Sevilla, Spain.

<sup>(3)</sup> Departamento de Sistemas Físicos, Químicos y Naturales. Universidad Pablo de Olavide, 41013 Sevilla, Spain.

<sup>(4)</sup> Department of Desertification and Geocology. Estación Experimental de Zonas Áridas, CSIC, 04120, Almería, Spain.

## Introduction

**Soil moisture** is a critical control of **evapotranspiration** or latent heat flux (LE) in dryland areas which cover 40% of the Earth. However, these areas also undergo periods of radiation-controlled evapotranspiration which compromises some satellite approaches estimating LE with surface temperature.

We tested the use of the **Temperature Vegetation Dryness Index (TVDI)** from Sandholt et al. (2002) to estimate the ratio of actual/potential LE using **MODIS 8-day satellite** data. We established the conditions for operational application in large regions with bioclimatic gradients and the accuracy of the method.

## Objectives


- Validate** surface energy fluxes derived from the *TVDI* at sites with different evapotranspiration controls and compare with alternative methods.
- Assess the role of **spatial heterogeneity** of climatic variables on *TVDI* errors.
- Sensitivity** of the *TVDI* to parameterizations for (i) Tair inputs, (ii) land cover types and (iii) algorithm determining hydrological boundaries

## Conclusion


- Validation:** *TVDI* was reliable to estimate latent and sensible heat fluxes only under *soil moisture-controlled evapotranspiration* conditions
- The effect of spatial heterogeneity in climatic variables depends also on LE controls.** Under energy-controlled LE, *TVDI* errors were related with radiation spatial heterogeneity. In soil moisture-controlled LE, errors were related with water vapor, while radiation was not significant.
- TVDI* accuracy was highly **sensitive** to the quality of Tair inputs and the algorithm used to extract hydrological boundaries.

## Methodology


**1. Regional estimates of *TVDI*** from triangle plots of radiometric temperature (*T<sub>s</sub>*) minus air temperature (*T<sub>air</sub>*) vs. *NDVI*. It **requires** *wet* and *dry* areas in each image and homogeneous climatic conditions in the region.



**Input data:** MODIS-Terra, meteorological station data in 2008




**2. Flux data from Eddy Covariance sites for validation of the *TVDI***




## Results

**a) Validation of surface fluxes from *TVDI*** at two field sites with Eddy Covariance data




**b) Role of spatial heterogeneity of climatic variables in Andalusia region on *TVDI* accuracy**



Correlation of *TVDI* errors with spatial CV of climatic variables in the region indicates an effect of spatial heterogeneity. E.g. the *TVDI* errors are not affected by spatial gradients in wind speed (WS) in water limited conditions.

**c) Sensitivity analysis of the *TVDI* to parameterizations**



Changes in correlation (R) and mean average error (MAE) of *TVDI* and the ratio (1-LE/LE<sub>p</sub>) from field data after changing different parameters.

Statistics during the year, GS (Grow season), and NGS (Non growing season).

**References in:** [Garcia, et al., 2014, Remote Sensing of Environment, Volume 149, 100-117](#)

**Acknowledgements:** Danish (FTP) Grant 09-070382, Andalusian regional government project Grant RNM-6685, the Spanish Ministry of Science and Innovation: CARBORAD (BBL2011-27493) FP7/2007-2013) grant agreement 244122..